



PROHEALTH BONE DENSITOMETRY

101- 45625 Hodgins Ave., Chilliwack, BC V2P 1P2
Phone 1 866 226 6143 Fax 1 866 236 8030
www.prohealthcr.ca

STANDARD OUT-PATIENT BONE DENSITOMETRY REQUISITION

ORDERING PRACTITIONER: ADDRESS, MSP PRACTITIONER NUMBER

Yellow highlighted fields must be completed to avoid delays in patient processing.

Consult provincial guidelines and protocols (www.BCGuidelines.ca) prior to completion

Bill to → MSP ICBC WorkSafeBC PATIENT OTHER: _____

PHN NUMBER		ICBC/WorkSafeBC NUMBER		LOCUM FOR PRACTITIONER:	
LAST NAME OF PATIENT		FIRST NAME OF PATIENT		MSP PRACTITIONER NUMBER	
DOB YYYY MM DD		SEX <input type="checkbox"/> M <input type="checkbox"/> F		PREGNANT <input type="checkbox"/> YES <input type="checkbox"/> NO	
PRIMARY CONTACT NUMBER OF PATIENT		SECONDARY CONTACT NUMBER OF PATIENT		OTHER CONTACT NUMBER OF PATIENT	
ADDRESS OF PATIENT		CITY/TOWN		PROVINCE	
DIAGNOSIS			CURRENT MEDICATIONS/DATE AND TIME OF LAST DOSE		

If this is a STAT order please provide contact telephone number:

Copy to Practitioner/MSP Practitioner Number/Address:

PERTINENT HISTORY - Follow-up examinations should be done at the same location (attach reports if available)

PREVIOUS BONE DENSITOMETRY <input type="checkbox"/> YES <input type="checkbox"/> NO	LOCATION	DATE
PREVIOUS LUMBAR SPINE X-RAYS <input type="checkbox"/> YES <input type="checkbox"/> NO	LOCATION	DATE

EXAMINATION REQUESTED

DIAGNOSTIC BONE MINERAL DENSITOMETRY (BMD)

BMD is only indicated if it is likely to alter treatment and is considered an MSP insured service for patients with moderate or higher risk of fracture, as outlined in the Osteoporosis Guideline at www.bcguidelines.ca. The risk can be determined using the FRAX calculator at www.shef.ac.uk/FRAX

- Example Risk Factors:
- Age > 65
 - Previous fragility fractures
 - Having a parent with fractured hip
 - Current smoking
 - Rheumatoid Arthritis
 - Secondary Osteoporosis
 - Alcohol consumption > 3 units/day
 - Glucocorticoids (≥ 7.5mg prednisone or equivalent daily for 3 months consecutively)

Check One:

- Moderate Risk (10 - 20% 10 year fracture risk) Recent Hip Fracture Hyperparathyroidism
 High Risk (>20% 10 year fracture risk) History of Fragility Fracture

FOLLOW-UP BMD MEASUREMENTS

There is insufficient evidence to recommend testing frequency for patients not taking OP medications. For patients on OP medications, repeat BMD exams are not justified based on current evidence and not considered medically necessary prior to **3 years** after the original measurement and only if it is likely to alter patient management.

- 3 or more years since prior BMD Exam Less than 3 years since BMD Exam (see below)

The following exceptions, as outlined in the Osteoporosis Guideline, may apply (check one):

- Patients receiving ≥ 7.5mg prednisone daily, or its equivalent for 3 months consecutively who require a baseline examination and repeat scans at 6 month intervals while on treatment.
 Patients in whom an early exam may be indicated: example - moderate and high risk patients on OP medications with multiple risk factors and test is likely to alter patient management.
 Primary Hyperparathyroidism
 Other specific high risk situations where repeat testing is likely to alter patient management

Specify _____

NON-DIAGNOSTIC BMD – PATIENT PAY

These are non-insured services for indications that are not covered by MSP, such as:

- Routine screening of men and women less than 65 years of age
- Part of routine screening around time of menopause
- Screening - the patient would like to proceed with the exam and pay privately
- Follow-up (when not clinically indicated) - the patient would like to proceed with the exam and pay privately
- Investigation of chronic back pain
- Investigation of exaggerated dorsal kyphosis

PATIENT HISTORY – please provide risk factors, therapies and other appropriate history			APPOINTMENT DATE AND TIME
TELEPHONE REQUISITION TIME	INITIALS OF RECORDER	DATE SIGNED (YYYY / MM / DD)	SIGNATURE OF REQUESTING PRACTITIONER

The personal information collected on this form is collected under the authority of the *Personal Information Protection Act*. The personal information is used to provide medical services requested on this requisition. The information collected is used for quality assurance management and disclosed to healthcare practitioners involved in providing care or when required by law. Personal information is protected from unauthorized use and disclosure in accordance with the *Personal Information Protection Act* and when applicable the *Freedom of Information and Protection of Privacy Act* and may be used and disclosed only as provided by those Acts.